

VII. Signal Enhancement By Tailing Of Reaction Products In The Invader™ Directed Cleavage Assay

It has been determined that when oligonucleotide probes are used in cleavage detection assays at elevated temperature, some fraction of the truncated probes will have been shortened by nonspecific thermal degradation, and that such breakage products can make the analysis of the target-specific cleavage data more difficult. Background cleavage such as this can, when not resolved from specific cleavage products, reduce the accuracy of quantitation of target nucleic acids based on the amount of accumulated product in a set timeframe. One means of distinguishing the specific from the nonspecific products is disclosed above, and is based on partitioning the products of these reactions by differences in the net charges carried by the different molecular species in the reaction. As was noted in that discussion, the thermal breakage products usually retain 3' phosphates after breakage, while the enzyme-cleaved products do not. The two negative charges on the phosphate facilitate charge-based partition of the products.

The absence of a 3' phosphate on the desired subset of the probe fragments may be used to advantage in enzymatic assays as well. Nucleic acid polymerases, both non-templated (*e.g.*, terminal deoxynucleotidyl transferase, polyA polymerase) and template-dependent (*e.g.*, Pol I-type DNA polymerases), require an available 3' hydroxyl by which to attach further nucleotides. This enzymatic selection of 3' end structure may be used as an effective means of partitioning specific from non-specific products.

In addition to the benefits of the partitioning described above, the addition of nucleotides to the end of the specific product of an invader-specific cleavage offers an opportunity to either add label to the products, to add capturable tails to facilitate solid-support based readout systems, or to do both of these things at the same time. Some possible embodiments of this concept are illustrated in Figure 67.

In Figure 67, an Invader™ cleavage structure comprising an Invader™ oligonucleotide containing a blocked or non-extendible 3' end (*e.g.*, a 3' dideoxynucleotide) and a probe oligonucleotide containing a blocked or non-extendable

3' end (the open circle at the 3' end of the oligonucleotides represents a non-extendible nucleotide) and a target nucleic acid is shown; the probe oligonucleotide may contain a 5' end label such as a biotin or a fluorescein (indicated by the stars) label (cleavage structures which employ a 5' biotin-labeled probe or a 5' fluorescein-labeled probe are shown below the large diagram of the cleavage structure to the left and the right, respectively). Following, cleavage of the probe (the site of cleavage is indicated by the large arrowhead), the cleaved biotin-labeled probe is extended using a template-independent polymerase (*e.g.*, TdT) and fluoresceinated nucleotide triphosphates. The fluorescein tailed cleaved probe molecule is then captured by binding via its 5' biotin label to streptavidin and the fluorescence is then measured. Alternatively, following, cleavage of a 5'-fluoresceinated probe, the cleaved probe is extended using a template-independent polymerase (*e.g.*, TdT) and dATP. The polyadenylated (A-tailed) cleaved probe molecule is then captured by binding via the polyA tail to oligo dT attached to a solid support.

The examples described in Figure 66 are based on the use of TdT to tail the specific products of Invader™-directed cleavage. The description of the use of this particular enzyme is presented by way of example and is not intended as a limitation (indeed, when probe oligos comprising RNA are employed, cleaved RNA probes may be extended using polyA polymerase). It is contemplated that an assay of this type could be configured to use a template-dependent polymerase, as described above. While this would require the presence of a suitable copy template distinct from the target nucleic acid, on which the truncated oligonucleotide could prime synthesis, it can be envisaged that a probe which before cleavage would be unextendible, due to either mismatch or modification of the 3' end, could be activated as a primer when cleaved by an invader directed cleavage. A template directed tailing reaction also has the advantage of allowing greater selection and control of the nucleotides incorporated.

The use of nontemplated tailing does not require the presence of any additional nucleic acids in the detection reaction, avoiding one step of assay development and troubleshooting. In addition, the use of non templated synthesis eliminated the step of hybridization, potentially speeding up the assay. Furthermore, the TdT enzyme is fast,

able to add at least >700 nucleotides to substrate oligonucleotides in a 15 minute reaction.

As mentioned above, the tails added can be used in a number of ways. It can be used as a straight-forward way of adding labeled moieties to the cleavage product to increase signal from each cleavage event. Such a reaction is depicted in the left side of Figure 66. The labeled moieties may be anything that can, when attached to a nucleotide, be added by the tailing enzyme, such as dye molecules, haptens such as digoxigenin, or other binding groups such as biotin.

In a preferred embodiment the assay includes a means of specifically capturing or partitioning the tailed invader-directed cleavage products in the mixture. It can be seen that target nucleic acids in the mixture may be tailed during the reaction. If a label is added, it is desirable to partition the tailed invader-directed cleavage products from these other labeled molecules to avoid background in the results. This is easily done if only the cleavage product is capable of being captured. For example, consider a cleavage assay of the present invention in which the probe used has a biotin on the 5' end and is blocked from extension on the 3' end, and in which a dye is added during tailing. Consider further that the products are to be captured onto a support via the biotin moiety, and the captured dye measured to assess the presence of the target nucleic acid. When the label is added by tailing, only the specifically cleaved probes will be labeled. The residual uncut probes can still bind in the final capture step, but they will not contribute to the signal. In the same reaction, nicks and cuts in the target nucleic acid may be tailed by the enzyme, and thus become dye labeled. In the final capture these labeled targets will not bind to the support and thus, though labeled, they will not contribute to the signal. If the final specific product is considered to consist of two portions, the probe-derived portion and the tail portion, can be seen from this discussion that it is particularly preferred that when the probe-derived portion is used for specific capture, whether by hybridization, biotin/streptavidin, or other method, that the label be associated with the tail portion. Conversely, if a label is attached to the probe-derived portion, then the tail portion may be made suitable for capture, as depicted on the right side of Figure 66. Tails may be captured in a number of ways,